

# Multiple Molecular H<sub>2</sub> Outflows in AFGL 618<sup>1</sup>

P. Cox<sup>2</sup>, P.J. Huggins<sup>3</sup>, J.-P. Maillard<sup>4</sup>, C. Muthu<sup>5</sup>, R. Bachiller<sup>5</sup>, T. Forveille<sup>6,7</sup>

## ABSTRACT

We report high spatial ( $0''.5$ ) and high spectral ( $9 \text{ km s}^{-1}$ ) resolution spectro-imaging of the  $2.12 \mu\text{m}$  H<sub>2</sub>  $1 \rightarrow 0$  S(1) line in the proto-planetary nebula AFGL 618 using *BEAR* at the CFHT. The observations reveal the presence of multiple, high-velocity, molecular outflows that align with the remarkable optical jets seen in *HST* images. The structure and kinematics of the outflows show how jets interact with circumstellar gas and shape the environment in which planetary nebulae form.

*Subject headings:* Planetary nebulae: general – Planetary nebulae: individual: AFGL 618 – Stars: AGB and post-AGB – Stars: circumstellar matter – Stars: winds and outflows

## 1. Introduction

High velocity ( $\gtrsim 100 \text{ km s}^{-1}$ ) directed outflows play a dramatic role in the rapid evolution of stars from the asymptotic giant branch (AGB) to the planetary nebula (PN) phase. The outflows interact with the roughly spherical molecular envelopes ejected by the stars on

---

<sup>1</sup>Based on observations collected at the Canada-France-Hawaii Telescope, operated by the National Research Council of Canada, the Centre National de la Recherche Scientifique de France, and the University of Hawaii.

<sup>2</sup>Institut d'Astrophysique Spatiale, Université de Paris Sud, F-91405 Orsay, France Pierre.Cox@ias.u-psud.fr

<sup>3</sup>Physics Department, New York University, 4 Washington Place, New York, NY 10003 patrick.huggins@nyu.edu

<sup>4</sup>Institut d'Astrophysique de Paris, C.N.R.S., 98b bd. Arago, F-75014 Paris, France maillard@iap.fr

<sup>5</sup>Observatorio Astronómico Nacional (IGN), Apt 1143, 28800 Alcalá de Henares, Spain muthu@oan.es bachiller@oan.es

<sup>6</sup>CFHT, PO Box 1597, Kamuela, HI 96743, USA forveill@cfht.hawaii.edu

<sup>7</sup>Observatoire de Grenoble, B.P. 53X, 38041 Grenoble Cedex, France

the AGB, producing strong shocks that pierce and disrupt the circumstellar gas (e.g., Cox et al. 2000; Huggins et al. 2000). Observations of shock diagnostics in objects that have only recently left the AGB are therefore of considerable interest in constraining this crucial evolutionary phase.

AFGL 618 is a bipolar proto-planetary nebula (PPN), which evolved from the AGB about 200 yr ago (Kwok & Bignell 1984). It was first identified by Westbrook et al. (1975) and since then it has been extensively observed. The presence of high-velocity molecular gas has been reported extending over  $\sim 250 \text{ km s}^{-1}$  in  $\text{H}_2 \text{ } 1 \rightarrow 0 \text{ S}(1)$  (Burton & Geballe 1986; Kastner et al. 2001) and  $\sim 400 \text{ km s}^{-1}$  in CO (Cernicharo et al. 1989; Gammie et al. 1989). High velocities are also seen in optical lines (e.g., Carsenty & Solf 1982) and the outflows appear as spectacular jet features in the *HST* images published by Trammell & Goodrich (2002). In this *Letter*, we report high velocity resolution ( $9 \text{ km s}^{-1}$ ) spectro-imaging of AFGL 618 in the  $\text{H}_2 \text{ } 1 \rightarrow 0 \text{ S}(1)$  line that reveals the detailed structure and kinematics of the shocked molecular gas, and its relation to the optical jets.

## 2. Observations

The observations were made on October 6, 2000 using the *BEAR* Imaging FTS at the f/35 infrared focus of the 3.60-meter Canada-France-Hawaii Telescope. A detailed description of *BEAR* and the data reduction is given by Maillard (2000). The  $256 \times 256$  HgCdTe facility camera provides a circular field of view of  $24''$  diameter, and the plate scale on the detector is  $0''.35/\text{pixel}$ . The seeing at  $2 \text{ } \mu\text{m}$  during the observations was typically  $0''.5$ . AFGL 618 was observed through a narrow band filter ( $4700 - 4750 \text{ cm}^{-1}$ ) which includes the  $\text{H}_2 \text{ } 1 \rightarrow 0 \text{ S}(1)$  line at  $4712.9 \text{ cm}^{-1}$  ( $2.12 \text{ } \mu\text{m}$ ). The raw data consist of a cube of 1150 planes with an integration time of 9 sec per image, an image being taken at each step of the interferometer. The maximum path difference corresponds to a resolution (FWHM) of  $0.137 \text{ cm}^{-1}$ , i.e.  $8.9 \text{ km s}^{-1}$ . A data cube of the star HD 162208 ( $m_K = 7.11$ ) was obtained for calibration. The  $2 \text{ } \mu\text{m}$  continuum emission of AFGL 618, which is found to be strongly peaked toward the center, has been subtracted from each plane of the cube. We have used the *HST* images and the J-band and  $\text{H}_2$  images of Ueta et al. (2001) to determine the co-ordinates of the *BEAR* data.

### 3. The Molecular Outflows

#### 3.1. Overview of the H<sub>2</sub> Emission

The velocity integrated image of AFGL 618 in the H<sub>2</sub> 1 → 0 S(1) line is shown in Fig. 1, together with the *HST* image in H $\alpha$ . The H<sub>2</sub> emission is dominated by two bipolar lobes which are oriented at PA  $\sim$  104°, and correspond to the jet system seen in the H $\alpha$  emission of the shocked-ionized gas.

The kinematic structure of the H<sub>2</sub> emission is found to be complex, and we present different aspects of the data in Figs. 2–5. The top panel in Fig. 2 shows the H<sub>2</sub> 1 → 0 S(1) spectrum of the whole nebula. The emission peaks at a velocity of  $-34.5 \pm 0.4$  km s<sup>-1</sup> (LSR), and shows broad wings extending from  $-200$  to  $+160$  km s<sup>-1</sup>, in agreement with previous observations (e.g., Burton & Geballe 1986). The peak is blue-shifted relative to the systemic velocity measured in CO ( $-21.8$  km s<sup>-1</sup>, Cernicharo et al. 1989), probably because of extinction. At low expansion velocities, maps of the H<sub>2</sub> are roughly symmetric about the center, but at higher velocities there is a clear blue/red separation to the east and west, respectively (Fig. 3). The nebula in H<sub>2</sub> is therefore tilted to the line of sight toward us in the east, consistent with the optical appearance.

The strongest H<sub>2</sub> emission lies  $\sim 2''$  each side of the center near the systemic velocity, but the emission near these positions also extends to the highest velocities. This is seen in spectra at these two positions (Fig. 2, center), which also show steep sided profiles in the red and blue wings to the east and west, respectively. The relation of these spectra to the overall kinematics is seen in the position-velocity diagram along the nebula axis shown in Fig. 4 – see also Kastner et al. (2001).

#### 3.2. Multiple Molecular Outflows

In addition to the main kinematic structure described above, a striking aspect of our data is the presence of localized, high-velocity features in the H<sub>2</sub> emission which are associated with the optical jets seen in the H $\alpha$  image. Fig. 1 already suggests some relation between the H<sub>2</sub> and H $\alpha$  emission, and this becomes evident in the H<sub>2</sub> channel maps at the extreme red- and blue-shifted velocities (Fig. 3). At these high velocities, discrete structures trace the locations of individual optical jets seen in the underlying *HST* image. These features can be followed in the data cube in position and velocity, and delineate the interaction of the jets with the molecular envelope.

On account of the tilt of AFGL 618 to the line of sight and the high optical extinction

toward the equatorial regions ( $A_V \gtrsim 10$  mag., e.g., Sánchez Contreras et al. 2002), there is less obscuration of the east lobe of the nebula. The optical jets are best seen on this side, and it is here that they show the clearest correspondence with the high velocity features seen in  $\text{H}_2$ . The correspondence is less complete in the west lobe, probably in part due to the higher optical extinction near the center, but also as a result of real differences (e.g., high velocity  $\text{H}_2$  lies part way along jet *a* in Fig. 3, but not along the whole length) which could reflect the geometry and/or strength of the interactions.

Fig. 5 presents a sequence of channel maps that show the relation between the optical jets and the  $\text{H}_2$  in the east lobe in detail. The origin of the optical jets cannot be traced near the star due to extinction, but  $\text{H}_2$  at intermediate velocities ( $-100 \text{ km s}^{-1}$ ) is detected near the base of the jets where they diverge from the center. At successively higher velocities, the  $\text{H}_2$  develops into separate, localized, structures that follow individual optical jets. Four separate structures can be seen corresponding to all four jets identified by Trammel & Goodrich (2002) in the east lobe (*b*, *c*, *d*, and *e*). Jet *b* is traced out to  $-163 \text{ km s}^{-1}$ . Jets *c* and *e* appear to be connected, with jet *c* terminating at  $-163 \text{ km s}^{-1}$ , and the associated jet *e* with an additional component extending to  $-178 \text{ km s}^{-1}$ . The highest velocities ( $-200 \text{ km s}^{-1}$ ) are associated with jet *d*, which is short and protrudes near the base of the jet system.

Three features of the observations provide further information on the relation of the jets to the molecular emission. First, although the  $\text{H}_2$  emission is not well spatially resolved across the jets, it is resolved along the extended jets *b* and *c+e*, and shows a velocity gradient, increasing away from the center. Second, unlike the  $\text{H}\alpha$ , the  $\text{H}_2$  emission is not strongly peaked at the head of the optical jets but extends downstream toward the star. Third, the  $\text{H}_2$  line widths observed near the heads of the jets are narrow ( $\approx 30 \text{ km s}^{-1}$ ) compared with the bulk velocity of the gas relative to the systemic velocity (Fig. 2, bottom panel).

## 4. Discussion

### 4.1. Geometry of the Outflows

The overall distribution and kinematics of the  $\text{H}_2$  suggests that the main emission arises in roughly formed, biconical cavities, consistent with the overall picture inferred from optical images. The steep-sided spectra (Fig. 2) and the sudden change from red- to blue-shifted gas near the systemic velocity (Fig. 3 & 4) indicate that the far side of the approaching (east) lobe and the near side of the receding (west) lobe lie close to the plane of the sky. The highest velocity gas then arises near the center, in the sides of the lobes that lie closest to the line of sight.

The east-west (blue-red) tilt of the optical jets (e.g., Sánchez Contreras et al. 2002) which form the lobes, and the angles of individual jets with respect to the line of sight estimated by Trammell & Goodrich (2002) ( $51^\circ$ ,  $31^\circ$ , and  $54^\circ$  for jets *b*, *c*, and *d*, respectively) are consistent with this general picture, although the short projected length of jet *d*, and the very high velocity that we observe in  $\text{H}_2$  suggest that it may be the closest to the line of sight. Given the short lifetime for optical emission from the jets (a few years, e.g., Sánchez-Contreras et al. 2002) compared to their dynamical time scales of  $\lesssim 200$  yr (see § 4.2), there could also be a history of outflows which partially excavated the biconical cavities in addition to those currently seen in the *HST* image.

## 4.2. Jet Interactions

Our observations cast light on the long-standing problem of the origin of the very high velocity  $\text{H}_2$  emission in AFGL 618 (e.g., Burton & Geballe 1986; Hartquist & Dyson 1987). Figs. 3 and 5 show that the emission arises from the interaction of jets with the circumstellar gas. The presence of  $\text{H}_2$  emission along the length of the optical jet structures is consistent with a class of models in which the ambient gas is shocked and dragged along with the jet through prompt or steady entrainment (e.g., Hartigan et al. 1996); similar cases are seen in the outflows around young stellar objects (e.g., Bachiller 1996). The kinematics that we observe place valuable constraints on such a picture. The  $\text{H}_2$  line widths near the heads of the jets ( $\approx 30 \text{ km s}^{-1}$ ) set upper limits on the turbulence which is believed to play a key role in the entrainment process. On the other hand, the bulk velocity of the molecular gas is extremely high. For example, the observed radial velocity of the  $\text{H}_2$  just behind the head of jet *b* is  $\sim 140 \text{ km s}^{-1}$  (relative to the systemic velocity); if the inclination to the line of sight is  $51^\circ$  (see above), the actual velocity is  $\sim 220 \text{ km s}^{-1}$ , and if the angle of inclination is  $66^\circ$  (Sánchez Contreras et al. 2002) the velocity is  $340 \text{ km s}^{-1}$ . For an adopted distance of 1.5 kpc to AFGL 618, the dynamical timescale of the jet is  $\lesssim 200$  yr.

The velocities are much too large for the molecular gas to survive acceleration in a single shock, for which the maximum velocity is  $\sim 45 \text{ km s}^{-1}$  (Draine, Roberge, & Dalgarno 1983). However, it is possible for the molecules to reform in the post-shock gas (e.g., Hollenbach & McKee 1989; Neufeld & Dalgarno 1989) although models of high velocity bow shocks including molecule reformation have not yet been developed.

It is striking that the  $\text{H}_2$  velocities that we observe are significantly higher than those indicated by observations of optical lines. The interpretation of optical line ratios at the heads of the jets using shock models yields shock velocities of only  $40\text{--}100 \text{ km s}^{-1}$  (Trammell & Goodrich 2002). The observed radial velocities in optical lines at the head of jet *b* are

$\sim 80 \text{ km s}^{-1}$  (Sánchez Contreras et al. 2002), and when corrected for an inclination of  $51^\circ$ – $66^\circ$  (see above) give values of  $\sim 130$ – $200 \text{ km s}^{-1}$ , the latter similar to shock velocities estimated from the optical line widths (Sánchez Contreras et al. 2002). However, independent of the inclination angle, the observed  $\text{H}_2$  velocity in jet *b* is a factor of  $\sim 1.8$  higher than the optical lines at the head.

If the  $\text{H}_2$  emission that we observe is shocked excited circumstellar gas, as seems most likely, then the high velocities that we observe suggest an even higher velocity, collimated driving wind. In addition, the differences between the  $\text{H}_2$  and optical line velocities indicate a range in shock conditions, which is not unlikely given the small scale structure seen in the  $\text{H}\alpha$  image. A possible, alternative scenario for the origin of the high velocity molecular gas is one in which the molecules are entrained or formed in the outflows close to their source, so are already present at high velocity and are excited by interactions with the ambient gas in the regions observed. Spectro-imaging of the optical lines with both higher spectral and spatial resolution than are currently available will be important to further study these issues.

#### 4.3. Evolution of the Nebula

The  $\text{H}_2$  observations reported here trace the detailed structure and kinematics of the molecular outflows in AFGL 618 which are also seen at lower angular resolution in other probes of the molecular gas. The full  $\text{H}_2$  spectrum shown in Fig. 2 resembles that of the CO lines (e.g., Cernicharo et al. 1989), and the shift from blue to red in going from the east to the west side of the nebula (Fig. 3) is also seen in millimeter molecular maps (e.g., Neri et al. 1992; Hajian et al. 1996; Meixner et al. 1998). However, the CO bullets reported by Ueta et al. (2001) at high negative velocities on the west side of the nebula are at variance with the overall kinematics reported here, and are probably not real.

The present observations provide evidence that the high velocity  $\text{H}_2$  outflows in AFGL 618 are the result of the interaction of jets with the circumstellar envelope. Similar examples of this type of interaction at different stages of development are seen in AFGL 2688 (Cox et al. 2000), M1-16 (Huggins et al. 2000), and NGC 7027 (Cox et al. 2002). All these cases support a view in which the geometry of the circumstellar gas around a newly forming PN is largely determined by the action of the jets, and plays a key role in determining the morphology of the fully formed PN.

We thank S. Trammell for providing a copy of the *HST*  $\text{H}\alpha$  image. We also thank A. Abergel, T. Ueta, and G. Pineau des Forêts for useful discussions, and an anonymous referee for helpful comments. This work has been supported in part by NSF grant AST-9986159

and a visiting professorship at the Université de Paris-Sud (to P.J.H.).

## REFERENCES

- Burton, M. G., & Geballe, T. R. 1986, MNRAS, 223, 13
- Bachiller, R. 1996, ARAA, 34, 111
- Carsenty, U., & Solf, J. 1982, A&A, 106, 307
- Cernicharo, J., Guélin, M., Martin-Pintado, J., Peñalver, J., & Mauesberger, R. 1989, A&A, 222, L1
- Cox, P., Lucas, R., Huggins, P. J., Forveille, T., Bachiller, R., Guilloteau, S., Maillard, J. -P., & Omont, A. 2000, A&A, 353, L25
- Cox, P., Huggins, P. J., Maillard, J. -P., Habart, E., Morisset, C., Bachiller, R., & Forveille, T. 2002, A&A, 384, 603
- Draine, B. T., Roberge, W. G., & Dalgarno, A. 1983, ApJ, 264, 485
- Gammie, C. F., Knapp, G. R., Young, K., Phillips, T. G., & Falgarone, E. 1989, ApJ, 345, L87
- Hajian, A. R., Phillips, J. A., & Terzian, Y. 1996, ApJ, 467, 341
- Hartigan, P., Carpenter, J. M., Dougados, C., & Skrutskie, M. F. 1996, AJ, 111, 1278
- Hartquist, T. W., & Dyson, J. E. 1987, MNRAS, 228, 957
- Hollenbach, D., & McKee, C. F. 1989, ApJ, 342, 306
- Huggins, P. J., Forveille, T., Bachiller, R., & Cox, P. 2000, ApJ, 544, 889
- Kastner, J. H., Weintraub, D. A., Gatley, I., & Henn, L., 2001, ApJ, 546, 279
- Kwok, S., & Bignell, R. C. 1984, ApJ, 276, 544
- Maillard, J. -P. 2000, in *Imaging the Universe in 3 Dimensions*, ed. E. van Breughel, & J. Bland-Hawthorn (ASP Conf. Ser. 195), 185
- Meixner, M., Campbell, M. T., Welch, W. J., & Likkell, L. 1998, ApJ, 509, 392
- Neri, R., García-Burillo, S., Guélin, M., Cernicharo, J., Guilloteau, S., & Lucas, R. 1992, A&A, 262, 544
- Neufeld, D. A., & Dalgarno, A. 1989, ApJ, 340, 869
- Sánchez Contreras, C., Sahai, R., & Gil de Paz, A. 2002, ApJ, 578, 269
- Trammell, S. R. & Goodrich, R. W. 2002, ApJ, 579, 688

Ueta, T, Fong, D., & Meixner, M. 2001, ApJ 557, L117

Westbrook, W. E., Willner, S. P., Merrill, K. M., Schmidt, M., Becklin, E., Neugebauer, G.,  
& Wynn-Williams, C. G. 1975, ApJ, 202, 407

Fig. 1.— Velocity integrated  $\text{H}_2$   $1 \rightarrow 0$  S(1) image of AFGL 618 observed with *BEAR* (bottom) and the *HST* image in  $\text{H}\alpha$  (top) from Trammell & Goodrich (2002). The  $\text{H}_2$  image is continuum subtracted, and the peak intensity is  $1.4 \times 10^{-5} \text{ W m}^{-2} \text{ sr}^{-1}$ . The coordinates of the field center are  $\alpha = 04^{\text{h}}42^{\text{m}}53^{\text{s}}.58$ ,  $\delta = 36^{\circ}06'53''.6$  (J2000.0).

Fig. 2.—  $\text{H}_2$   $1 \rightarrow 0$  S(1) spectra of AFGL 618. *Top*: integrated over the whole nebula. *Middle*:  $2''$  east and  $2''$  west (filled spectrum) of the center. *Bottom*: near tip of jet *b*. In the middle and bottom panels, the flux density is per pixel.

Fig. 3.—  $\text{H}_2$  channel maps of AFGL 618 superposed on the *HST*  $\text{H}\alpha$  image. *Top*: emission around the systemic velocity ( $-40$  to  $-3 \text{ km s}^{-1}$ ). *Middle*: blue and red intermediate velocities ( $-41$  to  $-125$  and  $-3$  to  $81 \text{ km s}^{-1}$ ). *Bottom*: blue and red extreme velocities ( $-126$  to  $-186$  and  $82$  to  $152 \text{ km s}^{-1}$ ). In the middle panel, the jets are labeled as in Trammell & Goodrich (2002). The contours are at 2, 6, and 10 to 100 by 20 percent of the peak intensity in each channel.

Fig. 4.— Position-velocity  $\text{H}_2$  map of AFGL 618, oriented at  $\text{PA} = 94^{\circ}.5$  (joining jets *a* and *b*, see Fig. 2). The contours are at 3.5, 7, 10.5, 14, and 20 to 100 by 10 percent of the peak intensity ( $10^{-5} \text{ W m}^{-2} \text{ sr}^{-1}$ ) of the map.

Fig. 5.— Sequence of  $\text{H}_2$  channel maps of the east lobe of AFGL 618 superposed on *HST*  $\text{H}\alpha$  image. The channel width is  $15 \text{ km s}^{-1}$  and the central velocity is given in each panel. The contours are 1, 1.5, 2, 3 to 30 by  $2.5 \times 10^{-7} \text{ W m}^{-2} \text{ sr}^{-1}$ . The jets are labeled as in Fig. 3.









